



## FluCov-Bulletin – June 2023

*FluCov project: combining data from around the world to better understand the impact of COVID-19 on influenza activity*

### Commentary

#### Contents

It is now more than three years since a cluster of atypical pneumonia cases in Wuhan, China, was reported to the World Health Organization (WHO) (January 1, 2020) that was later linked to the new **SARS-CoV-2** virus. The FluCov Bulletin provides an overview of the number of positive cases of **influenza** and **SARS-CoV-2** and the percentage of specimens that tested positive from January 2019 onwards in 22 countries across most regions of the world (see page 3).

#### Results

Globally, **influenza** activity has decreased (see Figure 1). The following country patterns were observed for **influenza** in June 2023:

- In the Southern Hemisphere, An increase in **influenza** detections was observed in **Australia** (mix of **influenza** A and B, not subtyped) and **South Africa** (**influenza** A(H3N2)), although the percentage of specimens that tested positive has started to decrease in **South Africa**.
- A smaller increase in **influenza** detections was also observed in **Brazil** (mix of **influenza** A(H1N1)pdm09 and B, lineage not determined), while the percentage of positive tests decreased.
- In the Northern Hemisphere, **influenza** activity is low in Europe (**United Kingdom, Poland, France, Germany, Netherlands, Spain**) and North America (**Canada, United States, Mexico**). In **Mexico**, the percentage of specimens that tested positive is increased for this time of the year compared to previous seasons.
- **Influenza** detections were low or decreasing in most Asian countries (**India, Japan, Philippines, Thailand** and **China**). A decrease in **influenza** detections was also observed in **South Korea**, following the increase in **influenza** A(H3N2), observed in May 2023.
- No update on **influenza** activity in May was available for **Israel, Italy** and **Vietnam**.

Globally, **SARS-CoV-2** detections have been generally low after the late 2022 peak in China (see Figure 1). The following country patterns were observed for **SARS-CoV-2** in June 2023:

- A small increase in **SARS-CoV-2** detections was observed in **Brazil** and **Thailand**.
- **SARS-CoV-2** detections were very low or decreasing, in most other countries in the Bulletin: **Australia, Canada, China, France, Germany, India, Israel, Italy, Mexico, Netherlands, Philippines, Poland, Spain, United Kingdom** and **Vietnam**.
- **SARS-CoV-2** detections also decreased in **South Korea**, however the number of detections remained high in June (over 400,000).
- No **SARS-CoV-2** detections were reported in **Egypt, Japan, South Africa** and the **United States**.

## Implications

In the Southern Hemisphere, an increase in **influenza** detections was observed in all countries covered by the Bulletin. The dominant virus type differed per country. In **South Africa** **influenza** A(H3N2) was dominant, while a mix of **influenza** A and B was reported in **Brazil** (A(H1N1)pdm09 and B) and **Australia** (influenza A subtype and B lineage not determined). An increase in **influenza** detections (A(H1N1)pdm09 if subtyped) was also observed in **Argentina**, while detections in **Chile** have started to decrease [1]. In the Northern hemisphere, influenza was generally low during the month of June, which is typical for the summer months.

The start of the **influenza** season in the Southern Hemisphere was slightly earlier (**South Africa**) or comparable to seasons before the COVID-19 pandemic (**Argentina** and **Chile** [2,3]). **Australia**, which had a particularly early **influenza** season in 2022 [4] is seeing a relatively late start of the season this year, compared to pre-pandemic seasons.

Thus far, only **influenza** B/Victoria was found to be the dominant lineage in countries covered by the Bulletin when this was determined. The detection and characterization of influenza B viruses has become increasingly important in the context of the COVID-19 pandemic, with the influenza B/Yamagata now appearing to be extinct [5].

Globally, **SARS-CoV-2** detections have been decreasing since the peak in **China** in December 2022 and were relatively low in June. In **South Korea** **SARS-CoV-2** activity was still high, detections peaked in May 2023 and started to decrease in June.

Importantly, WHO has announced that **SARS-CoV-2** is 'is now an established and ongoing health issue which no longer constitutes a public health emergency of international concern (PHEIC)' [6]. Despite this development, the surveillance of **SARS-CoV-2** will remain very important in the coming 12 months.




Figure 1: SARS-CoV-2 and influenza detections in the 22 countries covered by the Bulletin (period: from week 1/2019 to week 26/2023)

**Disclaimer: Comparisons between countries and seasons of influenza and SARS-CoV-2 detections should be made with care, as national surveillance systems may differ (e.g. surveillance structures and testing intensity) and change over time.**

## Monthly plots by country

The plots per country show weekly data for **influenza** and of **SARS-CoV-2** infections from 1 January, 2019 up to 2 July, 2023. This FluCov-Bulletin includes the countries Canada, United States, Mexico, Brazil, United Kingdom, France, Germany, Italy, Netherlands, Spain, Poland, South Africa, Egypt, China, Japan, South Korea, India, Philippines, Thailand, Vietnam, Israel and Australia.

Per country, the first plot displays the number of positive **influenza** (in blue) and **SARS-CoV-2** (in red) detections. An overview of the absolute number of **influenza** and of **SARS-CoV-2** detections per country can be found on [pages 26-28 of this FluCov-Bulletin \(click here\)](#). The bar displays the Stringency Index (SI; a country-specific composite metric of the mitigation measures that are in place) over time. The second plot shows the **influenza** detections by subtypes/lineages reported to FluNet. The third plot displays the percentage of specimens testing positive for **influenza** during the current season (in red), the last season, and the average of the two pre COVID-19 seasons (2017-18 and 2018-19).

### The FluCov Dashboard is live!

All Figures and Tables in the FluCov-Bulletin can now be accessed (real-time) at:

<https://www.nivel.nl/en/dossier-epidemiology-respiratory-viruses/flu-cov-dashboard>

### Countries (click to view plot)

North America

Canada

United States

Central America Caribbean

Mexico

Tropical South America

Brazil

Northern Europe

United Kingdom

Eastern Europe

Poland

South West Europe

France

Germany

Italy

Netherlands

Spain

Northern Africa

Egypt

Southern Africa

South Africa

Eastern Asia

China

Japan

South Korea

Southern Asia

India

South East Asia

Philippines

Thailand

Vietnam

Western Asia


Israel

Oceania


Australia

# North America


## Canada




## Percentage of specimens testing positive for influenza in different seasons



## United States




## Percentage of specimens testing positive for influenza in different seasons




# Central America Caribbean

## Mexico




## Percentage of specimens testing positive for influenza in different seasons




# Tropical South America

## Brazil




## Percentage of specimens testing positive for influenza in different seasons



# Northern Europe

## United Kingdom




**Percentage of specimens testing positive for influenza in different seasons: data not available**




# Eastern Europe

## Poland




## Percentage of specimens testing positive for influenza in different seasons




# South West Europe


## France




## Percentage of specimens testing positive for influenza in different seasons




## Germany




## Percentage of specimens testing positive for influenza in different seasons




## Italy



## Percentage of specimens testing positive for influenza in different seasons




## Netherlands




**Percentage of specimens testing positive for influenza in different seasons: data not available**

## Spain




## Percentage of specimens testing positive for influenza in different seasons




# Northern Africa

## Egypt




## Percentage of specimens testing positive for influenza in different seasons




# Southern Africa

## South Africa




## Percentage of specimens testing positive for influenza in different seasons






# Eastern Asia

## China




Note: Due to a high number of retrospectively added SARS-CoV-2 detections, the peak in China in late 2022 increased significantly, compared to previous Bulletins.

## Percentage of specimens testing positive for influenza in different seasons




## Japan




***Percentage of specimens testing positive for influenza in different seasons: data not available***

## South Korea




## Percentage of specimens testing positive for influenza in different seasons




# Southern Asia

## India




## Percentage of specimens testing positive for influenza in different seasons




## South-East Asia

### Philippines




**Percentage of specimens testing positive for influenza in different seasons: data not available**


## Thailand



## Percentage of specimens testing positive for influenza in different seasons




## Vietnam




**Percentage of specimens testing positive for influenza in different seasons: data not available**

# Western Asia

## Israel




## Percentage of specimens testing positive for influenza in different seasons






# Oceania

## Australia



## Percentage of specimens testing positive for influenza in different seasons



## Absolute numbers per country

Country	Year	Cases <sup>a,b</sup> of SARS-CoV-2	+/- since last month <sup>c</sup>	Cases <sup>a</sup> of influenza	+/- since last month <sup>c</sup>	Week of last influenza update
Australia	2019			12,404		
Australia	2020	28,381		784		
Australia	2021	338,226		8		
Australia	2022	10,418,952		12,312		
Australia	2023	723,275	85,167	3,553	2,087	2023-26
Brazil	2019			3,320		
Brazil	2020	7,563,551		1,361		
Brazil	2021	14,700,856		1,240		
Brazil	2022	14,038,581		3,648		
Brazil	2023	1,353,635	77,022	18,322	2,416	2023-26
Canada	2019			43,196		
Canada	2020	565,508		44,956		
Canada	2021	1,536,966		337		
Canada	2022	2,390,153		71,314		
Canada	2023	196,090	10,480	10,743	758	2023-26
China	2019			122,757		
China	2020	96,673		31,164		
China	2021	35,398		25,812		
China	2022	86,958,455		56,455		
China	2023	14,364,114	19,645	118,792	839	2023-26
Egypt	2019			1,998		
Egypt	2020	136,644		659		
Egypt	2021	248,084		233		
Egypt	2022	130,805		2,709		
Egypt	2023	490	0	594	20	2023-22
France	2019			25,405		
France	2020	2,564,972		16,589		
France	2021	6,917,610		3,071		
France	2022	28,717,231		40,126		
France	2023	999,835	45,306	18,759	68	2023-26
Germany	2019			1,215		
Germany	2020	1,734,470		958		
Germany	2021	5,430,510		29		
Germany	2022	30,220,061		1,923		
Germany	2023	1,194,476	8,523	539	4	2023-26
India	2019			10,426		
India	2020	10,266,679		652		
India	2021	24,572,130		5,093		
India	2022	9,840,329		1,663		
India	2023	315,804	3,600	1,491	233	2023-26
Israel	2019			1,796		
Israel	2020	419,661		1,424		
Israel	2021	962,277		456		
Israel	2022	3,381,643		774		
Israel	2023	65,537	2718	801	0	2023-13

Country	Year	Cases <sup>a,b</sup> of SARS-CoV-2	+/- since last month <sup>c</sup>	Cases <sup>a</sup> of influenza	+/- since last month <sup>c</sup>	Week of last influenza update
Italy	2019			2,787		
Italy	2020	2,083,689		7,485		
Italy	2021	3,897,739		31		
Italy	2022	19,187,010		5,817		
Italy	2023	729,363	30,849	2,415	0	2023-17
Japan	2019			10,343		
Japan	2020	230,304		2,915		
Japan	2021	1,503,484		9		
Japan	2022	27,478,747		258		
Japan	2023	4,698,502	0	2,670	50	2023-25
Mexico	2019			6,963		
Mexico	2020	1,496,067		4,799		
Mexico	2021	2,538,755		960		
Mexico	2022	3,236,805		10,314		
Mexico	2023	361,713	11,923	2,488	401	2023-26
Netherlands	2019			5,145		
Netherlands	2020	785,874		3,168		
Netherlands	2021	2,329,020		471		
Netherlands	2022	5,454,287		11,082		
Netherlands	2023	50,108	701	9,191	31	2023-25
Philippines	2019			612		
Philippines	2020	472,523		52		
Philippines	2021	2,371,346		105		
Philippines	2022	1,218,790		260		
Philippines	2023	101,701	22,825	212	85	2023-26
Poland	2019			1,786		
Poland	2020	1,297,400		1,282		
Poland	2021	2,811,801		2		
Poland	2022	2,259,187		1,604		
Poland	2023	149,504	940	1,864	3	2023-26
South Africa	2019			1,164		
South Africa	2020	1,039,161		157		
South Africa	2021	2,407,371		413		
South Africa	2022	602,048		1,171		
South Africa	2023	23,953	0	847	330	2023-26
South Korea	2019			1,702		
South Korea	2020	60,722		505		
South Korea	2021	574,528		0		
South Korea	2022	28,481,550		295		
South Korea	2023	3,072,333	428,095	729	111	2023-26
Spain	2019			16,580		
Spain	2020	1,955,216		8,827		
Spain	2021	4,550,685		2,206		
Spain	2022	7,178,335		18,099		
Spain	2023	215,776	18,082	8,736	74	2023-26
Thailand	2019			1,568		
Thailand	2020	6,919		297		
Thailand	2021	2,216,551		23		
Thailand	2022	2,500,484		575		
Thailand	2023	27,644	9,605	385	24	2023-24

Country	Year	Cases <sup>a,b</sup> of SARS-CoV-2	+/- since last month <sup>c</sup>	Cases <sup>a</sup> of influenza	+/- since last month <sup>c</sup>	Week of last influenza update
United Kingdom	2019			42,447		
United Kingdom	2020	2,563,565		14,377		
United Kingdom	2021	10,878,110		2,755		
United Kingdom	2022	10,752,351		26,905		
United Kingdom	2023	444,641	15,435	5,623	144	2023-26
United States	2019			268,524		
United States	2020	19,577,585		229,766		
United States	2021	33,956,701		39,507		
United States	2022	45,877,410		470,074		
United States	2023	4,025,133	0	42,242	2,280	2023-26
Vietnam	2019			355		
Vietnam	2020	1,456		146		
Vietnam	2021	1,729,801		39		
Vietnam	2022	9,793,974		106		
Vietnam	2023	95,338	9,106	235	0	2023-20

<sup>a</sup> Laboratory-confirmed cases.

<sup>b</sup> As of the 24<sup>th</sup> bulletin, the data source, used by Our World In Data, for SARS-CoV-2 cases has been changed retrospectively. As a result, yearly totals displayed in this table may differ from those in previous bulletins.

<sup>c</sup> Influenza cases are reported by FluNet on a weekly basis. To convert these data to months, weekly data are assigned to the month most days in that week belong to. SARS-CoV-2 cases are reported per day and assigned to each month by date. +/- since last month includes all cases over the last full calendar month.

# Methodology

## Background

After assessment of alarming levels of spread and severity of SARS-CoV-2 virus, on March 11, 2020, WHO declared COVID-19 a pandemic [7]. The emergence of this new virus has had a major impact on the global circulation of respiratory viruses, including influenza and RSV [8]. The FluCov project aims to understand and communicate the impact of COVID-19 on: i) influenza activity and ii) prevention and control measures (e.g. vaccination) in the coming years.

## Scope

The countries included in this FluCov-Bulletin are distributed over the Americas (North, Central and Tropical South), Europe (Northern, South West and Eastern), Africa (Northern and Southern), Asia (Eastern, Southern, South East and Western) and Oceania. These data are compared to the prevention and control measures applied in each country using the Stringency Index from the Oxford COVID-19 Government Response Tracker (OxCGRT) [9].

## Data sources

- **Influenza:** FluNet [10] is a global web-based tool for influenza virological surveillance first launched in 1997. The virological data entered into FluNet, e.g. number of influenza viruses detected by subtype, are critical for tracking the movement of viruses globally and interpreting the epidemiological data. The data are provided remotely by National Influenza Centres (NICs) of the Global Influenza Surveillance and Response System (GISRS) and other national influenza reference laboratories collaborating actively with GISRS or are uploaded from WHO regional databases.
- **SARS-CoV-2:** Our World in Data systematically collects COVID-19 data which is presented in their online tool [11]. We used this platform to extract data on the number of cases, as well as tests performed per country. As of 8 March 2023, Our World in Data changed their primary data source from the John Hopkins repository on daily confirmed COVID-19 cases to the WHO [12].
- **Government response tracker:** The Oxford COVID-19 Government Response Tracker (OxCGRT) [9] systematically collects information on several different common policy responses that governments have taken to respond to the pandemic on 20 indicators such as school closures and travel restrictions. It now has data from more than 180 countries. OxCGRT data is downloaded directly from the Our World in Data platform.

## Extraction details

Data were extracted on 10 July 2023 and cover the period 1 January 2019 to 2 July 2023 (influenza) and 6 July (SARS-CoV-2). Data from both platforms are regularly updated and **sometimes retrospectively corrected**. This might explain any discrepancies between our reported figures and the data published online, even when using data for the exact same period. In case of any unclarities or perceived irregularities, feel free to contact us at [flu cov@nivel.nl](mailto:flu cov@nivel.nl).

## References

- [1] PAHO. Influenza Situation Report. Weekly Summary: Epidemiological Week 25 2023. <https://www.paho.org/en/influenza-situation-report> [accessed 11 July 2023]
- [2] WHO. FluNet. <https://www.who.int/tools/flunet> [accessed 11 July 2023]
- [3] Ministerio de Salud Argentina. Comunicación epidemiológica: Circulación de Influenza, VSR y otros virus respiratorios. SE23/2023. 9 June 2023. <comunicacion-influenza-ovr-06-2023.pdf> (salud.gob.ar) [accessed 11 July 2023]
- [4] Nivel. FluCov-Bulletin – May 2022. Utrecht; 2022. Report No.: 10.
- [5] Paget J, Caini S, Del Riccio M, van Waarden W, Meijer A. Has influenza B/Yamagata become extinct and what implications might this have for quadrivalent influenza vaccines? Euro Surveill. 2022 Sep;27(39):2200753. doi: 10.2807/1560-7917.ES.2022.27.39.2200753.
- [6] WHO. Statement on the fifteenth meeting of the IHR (2005) Emergency Committee on the COVID-19 pandemic. 5 May 2023. [Statement on the fifteenth meeting of the IHR \(2005\) Emergency Committee on the COVID-19 pandemic \(who.int\)](Statement on the fifteenth meeting of the IHR (2005) Emergency Committee on the COVID-19 pandemic (who.int)) [accessed 9 May 2023]
- [7] WHO. Listing of WHO's response to COVID-19. <https://bit.ly/3mIMtRi> [accessed 1 July 2022]
- [8] WHO. Influenza Update N° 416. <http://bit.ly/3T5SvHV> [accessed 7 April 2022]
- [9] Oxford COVID-19 Government Response Tracker, Blavatnik School of Government, University of Oxford. <http://bit.ly/41WqmQX> [accessed 16 June 2021]
- [10] WHO. FluNet. <https://www.who.int/tools/flunet> [accessed 8 March 2023]
- [11] Ritchie, H., Ortiz-Ospina, E., Beltekian, D., Mathieu, E., Hasell J., Macdonald B. et al. Coronavirus Pandemic (COVID-19). <https://ourworldindata.org/coronavirus> [accessed 15 June 2021]
- [12] Mathieu E, Rodés-Guirao L. Our World in Data will rely on data from the WHO to track confirmed COVID-19 cases and deaths. <https://ourworldindata.org/covid-jhu-who> [accessed 5 April 2023]

---

### Project Team

**Nivel, Netherlands:** Bronke Boudewijns, Marco Del Riccio, Willemijn van Waarden, Saverio Caini, John Paget

#### Global Influenza Initiative:

Ben Cowling, School of Public Health, University of Hong Kong, Hong Kong

Ann Falsey, Rochester General Hospital, University of Rochester School of Medicine, Rochester, NY

Angele Gentile, Ricardo Gutiérrez Children's Hospital, Buenos Aires

Jan Kyncl, Department of Infectious Diseases Epidemiology, National Institute of Public Health, Prague

Bruno Lina: Virpath Laboratory, University of Lyon, Lyon

Raina McIntyre: The Kirby Institute, University of New South Wales, Sydney



Global **Influenza** Initiative

**Sanofi, France:** Erica Dueger, Clotilde El Guerche-Séblain, Meral Akçay, Cecile Eymin

### Websites

Project Website: <https://www.nivel.nl/en/fluov>

FluCoV Dashboard: <https://www.nivel.nl/en/dossier-epidemiology-respiratory-viruses/fluov-dashboard>

### Funding

The FluCov Project is funded by Sanofi, France.

---